

## BY 3 DAYS OF AGE:

SUPPLEMENTATION with IRON:

**Ferroforte** (comes in 100mL bottle) – **1mL DOSE**

- **Piglets:**
  - ALL piglets to get one dose no later than 3 days of age



## AT WEANING:

VACCINATION against WASTING DISEASES:

*Circo Virus*

**Circoflex** (comes in 50 dose bottles) – **1mL DOSE**

**21 day withdrawal**

- **Piglets:**
  - One dose ONCE at 3 weeks of age



## AT WEANING AND 3-4 WEEKS LATER:

VACCINATION against SOIL-BORNE disease:

*Erysipelas*

**ER Bac Plus** (comes in 50 dose bottles) – **2mL DOSE**

**21 day withdrawal**

- **Piglets:**
  - 2ml dose at >4 weeks of age
  - Booster 2ml dose 3-4 weeks after first dose



VACCINATION against SOIL-BORNE disease:

*Clostridial Disease*

**Ultrachoice 8** (comes in 50 dose bottles) – **2mL DOSE**

**21 day withdrawal**

- **Piglets:**
  - 2ml dose at weaning
  - Booster 2ml dose 3-4 weeks after first dose



For more information on best practices for raising pigs outdoors, refer to the *Canadian Small-Scale Pig Farming Manual*: [www.casv-acvp.com/small-scale-pig-farming.html](http://www.casv-acvp.com/small-scale-pig-farming.html)

## AROUND 6-10 WEEKS OF AGE AND 4-5 WEEKS BEFORE SLAUGHTER:

### DEWORMING:

**Safeguard** (does not treat mites or threadworms)

**0 day withdrawal**

OR

**Ivomec** (does not treat trichuris suis, but does treat mange/mites)

**28 day withdrawal (injectable)**

**14 day withdrawal (top dress)**

- **Finishing Pigs:**

- Treat between 6 and 10 weeks old
- Treat 4-5 weeks prior to slaughter (about 21 to 25 weeks)



***If you find milk spot livers after one turn of this deworming protocol, add in a 3<sup>rd</sup> deworming dose in the finisher pigs (about week 15/16)!!!***

***\*\*\*Note: you can rotate dewormers\*\*\****

